

# **ANTIDIABETIC POTENTIAL OF ABELMOSCHUS ESCULENTUS LINN. IN ALLOXAN- INDUCED DIABETIC RATS**

**Authors: Prabhat Jain**

Scan Research Laboratories,  
Bhopal, Madhya Pradesh,  
India – 462041

**Nilesh Jain**

Sagar Institute of Research Technology &  
Science-Pharmacy, Ayodhya Bypass RoadBhopal,  
Madhya Pradesh, India – 462041

**Geeta Parkhe**

Scan Research Laboratories,  
Bhopal, Madhya Pradesh,  
India – 462041

**Ruchi Jain**

Sagar Institute of Research Technology &  
Science-Pharmacy, Ayodhya Bypass RoadBhopal,  
Madhya Pradesh, India – 462041

**Abstract**

**Abelmoschus esculentus Linn. (Malvaceae) is described in the folk medicine for the treatment of diabetes mellitus. The monech seeds of Abelmoschus esculentus are having potential in the development of drug for diabetes due to their antidiabetic activity. The hypoglycemic effect of the extract was tested in normal, glucose loading and alloxan-induced diabetic rats. Aqueous and ethanolic extracts (250 and 500 mg/kg body weight), were administered orally to male Wistar albino rats. Total phenolic content was estimated in the extracts. The parameters studied included oral glucose tolerance test, fasting blood glucose, serum insulin and glycated haemoglobin levels, liver glycogen content, serum lipid profile, and changes in body weights. The extracts produced a dose-dependent fall in fasting blood glucose (FBG). After 15 days of treatment with extracts the maximum reduction in FBG (35.14%) was observed in diabetic rats treated with ethanolic**

**extract 500 mg/kg dose. Serum lipid levels were reversed towards near normal and a control in the loss of body weight was observed in treated rats as compared to diabetic control. The extract treatment also showed a significant increase in the liver glycogen and a significant decrease in glycated haemoglobin levels.**

**Keywords**

**Abelmoschus esculentus Linn, Antidiabetic activity, oral glucose tolerance test, Aqueous and ethanolic extracts, Insulin,**

**Introduction**

Diabetes mellitus is a metabolic disorder characterized by hyperglycemia and alterations in carbohydrate, fat and protein metabolism, associated with absolute or relative deficiencies in insulin secretion and/or insulin action. Though different types of oral hypoglycemic agents are available along with insulin for the treatment of diabetes mellitus, there is a growing interest in herbal remedies, due to the side effects associated with these therapeutic agents. Because of

perceived effectiveness, minimal side effects in clinical experience and relatively low cost, herbal drugs are widely prescribed even when their biologically active compounds are unknown<sup>1</sup>.

Synthetic hypoglycemic agents can produce serious side effects and in addition, they are not suitable for use during pregnancy<sup>2</sup>. Therefore the search for more effective and safer hypoglycemic agents has continued to be an important area of active research. Furthermore, after the recommendations made by WHO on diabetes mellitus<sup>3</sup>, investigations on hypoglycemic agents from medicinal plants have become more important.

*Abelmoschus esculentus* (Linn.) Moench (Family:- Malvaceae) is an annual or perennial herb, growing to 2 m tall. The leaves are heart shaped, 10–20 cm long and broad, palmately lobed with 5–7 lobes. The large, yellow, hibiscus-like flowers are 4–8 cm diameter, with five white to yellow petals, often with a red or purple spot at the base of each petal. The seed pods are 3 to 10 inches long, tapering, usually with ribs down its length. These tender, unripe seed pods are used as a vegetable, and have a unique texture and sweet flavor. The pods, when cut, exude a mucilaginous juice that is used to thicken stews (gumbo), and have a flavor somewhat like a cross between asparagus and eggplant. The fruit is a capsule up to 18 cm long, containing numerous seeds. Its also known as Lady's Fingers, gumbo, gumbo, ochro, bamia, bamie, quiabo. In Spanish okra is quibombo; the French word is gombo, bamia or banya, in India it is bhindi, and in the eastern Mediterranean and Arab countries bamies<sup>4</sup>.

The fruit is highly proteinaceous. It is a good source of vitamin A and vitamin C. It is low in calories and is fat-free. It has also considerable medicinal and

industrial value<sup>5</sup>. A mucilagenous preparation from the pod can be used as a plasma replacement<sup>6,7</sup>.

Okra mucilage has medicinal applications; when used as a plasma replacement or 'blood-volume expander'. The mucilage of Okra not only binds cholesterol but the bile acid carrying toxins dumped into it by the filtering liver. It also has industrial applications; when added as size to glaze paper and used in confectionary<sup>8-10</sup>.

Okra constitutes minerals, vitamins, proteins, carbohydrates, enzymes and very high quantities of mucilages<sup>11</sup>. *Abelmoschus esculentus* L. (or *Hibiscus esculentus* or Okra) – Malvaceae is used for a long time as an edible vegetable in many countries, and commonly eaten in Vietnam because of its nourishing components. Traditionally, it is believed that the plant is useful in the treatment of inflammatory disorders, constipation, retention of urine, On the other hand, a number of previous studies have reported that *Abelmoschus* sp. possessed hypoglycemic effect. However, there is a little study regarding its hypolipidemic effect.

Fresh fruits are cut into small pieces and then boiled in water, to obtain mucilaginous decoction. This decoction is prescribed for cough, throat infection and bronchitis. Wehmer records that the fruit contains abundant pectin; mucilage; starch; some fat, 4 percent; water 80.7 percent; and ash, 1.41 percent. Popp analyzed the seeds and found nitrogen, 2.4 to 2.5 percent; their ash finding K<sub>2</sub>O 39 percent; MgO 12 percent, CaO 7.8 percent and P<sub>2</sub>O<sub>5</sub> 24.7 percent. Jamieson and Baugham analyzed the seeds of okra; their results are as follows: palmitic acid, 27.23 percent; stearic acid, 2.75 percent; arachidic acid, 0.05 percent; oleic acid, 43.74 percent; linolic acid, 26.62 percent; Unsaponifiable matter, 0.37 percent. Read

reports that the roots contain gum, 16 percent; and the seeds, vitamin C<sup>12-17</sup>.

## **Material and Method**

### **Collection of plant material**

The species for the proposed study that is *Abelmoschus esculentus* Linn., Fruits were collected, from the local market of bhopal (M.P.) India were authenticated by Agriculture College, Indore (M.P.) with the help of forest department botanist.

### **Preparation of the plant extracts**

The fruits were washed properly with water to remove the mud or dust if any. Initially it was dried in a sun for an hour than was shade dried completely. The dried fruit were then powdered by means of wood grinder and was sieved through sieve no. 40 and then stored in airtight containers. The powdered fruit (150 g) were extracted with ethanol (90% w/v) for 24 h using a soxhlet extractor. The powdered fruits (30 g) were extracted with distilled water by cold maceration method. This extracts were concentrated to dryness under reduced pressure and controlled temperature (50-60°C) to yield solid masses that were completely free from solvents (yield: 18.92% w/w and 12.36 %W/W for aqueous extract and ethanolic extract respectively). The animals treated with 250 and 500 mg/kg body weight of the Ethanolic and aqueous extracts which were suspended in 0.5% Tween 80 in saline (0.9% w/v) for oral administration.

### **Animals**

Animal ethical clearance was obtained from Ethics Committee (1429/PO/a/11/CPCSEA) of Sagar Institute of Research & technology-Pharmacy, Bhopal, India Healthy Male wistar albino rats (100-200 gm) were selected for the present investigation. The rats were housed in a well-ventilated, temperature-controlled (27 ± 2°C, RH 60-70%) animal room with 12-12 hr light dark cycle for 7 days prior to the

experimental period. The animals were provided with standard pellet diet and water ad. Libitum. in polypropylene cages. Animals were periodically weighed before and after experiments.

### **Experimental Procedures**

#### **Oral glucose tolerance test**

The oral glucose tolerance test (OGTT) was performed on overnight fasting (12 h) normal rats. Vehicle, Ethanolic and aqueous extract of *Abelmoschus esculentus* fruits (250 and 500 mg/kg, per os) and glibenclamide (2.5 mg/kg b.w. per os) were administered to six groups of rats, respectively. Glucose (4 g/kg, per os) was fed 90 min after pretreatment with vehicle, Ethanolic and aqueous extract of *Abelmoschus esculentus* or glibenclamide. Blood sample (0.3 ml) was withdrawn from the retro-orbital plexus under ether anesthesia at 0, 30, 60, 90 and 120 min of extract or glibenclamide administration. At the end of each experiment, polyvidone iodine solution (Betadine®) was applied at the site of the retro-orbital puncture to prevent infection. blood glucose levels were determined using a glucometer<sup>18,19</sup>.

#### **Study on diabetic rats (non-insulin dependent diabetes model—NIDDM)**

##### **Induction of diabetes**

Diabetes was induced in rats by the single intraperitoneal (i.p.) injection of of freshly prepared solution of alloxan monohydrate at a dose of 150 mg/kg b.w. dissolved in distilled water (1 ml/kg b.w.). Seven days after the injection, the blood glucose levels were measured. Each animal with a blood glucose concentration level above 250g/dl was considered to be diabetic and used in the experiments<sup>18-21</sup>. To prevent the hypoglycemia, which occurred during the first 24 h following the alloxan administration, 5% glucose solution was orally given to the diabetic rats.

In all experiments, rats were fasted for 16 h prior to alloxan monohydrate injection.

Fourty two animals were divided into seven groups of six animals each as follows:

Group 1 -: Served as normal control (0.9 % w/v Saline 10 ml/kg b.w./day)

Group 2 -: Served as diabetic control (Alloxan induced)

Group 3 -: Received aqueous extract, 250 mg/kg b.w. /day orally

Group 4 -: Received Ethanolic extract, 250 mg/kg b.w. /day orally

Group 5 -: Received aqueous extract, 500 mg/kg b.w. /day orally

Group 6 -: Received Ethanolic extract, 500 mg/kg b.w. /day orally

Group 7 -: Served as reference standards (Glibenclamide, 2.5 mg/kg b.w. /day)

#### **Acute anti diabetes effect of test samples**

The test samples (Ethanolic extracts, aqueous extracts, and glibinclinamide) were administered orally by using a gastric gavage needle. The blood sample was collected from rats by retro orbital plexus bleeding method and blood glucose levels were determined at 0, 2, 4, 6 and 24 hour after the single oral administration of the test samples by using glucometer<sup>22</sup>.

#### **Subacute anti diabetes effect of test samples**

The test samples (ethanolic extracts, aqueous extracts, and glibinclinamide) were administered orally for 14 days consecutively. Blood glucose levels were determined at 10:00 a.m. on 1st, 3rd, 5th and 14th days after the single oral administration of the test samples by using glucometer<sup>22</sup>.

#### **Statistical analysis**

Values are presented as means  $\pm$  S.E.M. Statistical differences between the treatments and the controls were tested by one-way analysis of variance

(ANOVA) followed by the Dunnett test using the “SYSAT” statistic computer program. A difference in the mean values of  $P < 0.01$  or less was considered as to be statistically significant.

#### **Results**

The phytochemical screening of *Abelmoschus esculentus* revealed the presence of Alkaloids, Carbohydrates, Terpenoids and Flavonides.

#### **Glucose tolerance test**

The effect of the different doses of aqueous and Ethanolic extract of *Abelmoschus esculentus* on the fasting blood glucose levels of both normal and treated rats are given in Table 1. The result showed the blood glucose level had decreased significantly by 60 min (with respect to 30 min level) and this was maintained until 120 min. The Ethanolic extract at the dose of 500 mg/kg showed a significant reduction ( $P < 0.01$ ) in blood glucose level within 120 min.

#### **Antidiabetic activity**

The single administration of (acute study) Ethanolic extract (500 mg/kg b.w.) has more significantly reduced the blood glucose level at 4<sup>th</sup> hrs from  $130.14 \pm 1.4$  to  $113.23 \pm 0.03$  and significant hypoglycemia was maintained up to 24<sup>th</sup> hrs. Glibenclamide (2.5 mg/kg) has also significantly reduced the blood glucose level at 4<sup>th</sup> hrs  $131.00 \pm 1.61$  to  $106.33 \pm 0.92$  and significant hypoglycemia was maintained up to 24<sup>th</sup> hr table-2. On repeated administration (subacute study) vehicle, glibenclamide, aqueous and Ethanolic extract of *Abelmoschus esculentus* for 14 days, a significant ( $P < 0.01$ ) decrease in blood glucose level of the diabetic rats were seen at a dose of (250, 500 mg/kg b.w.), in dose dependent manner as compared to diabetic control group. On the other hand glibenclamide showed a significant ( $P < 0.01$ ) decrease in blood glucose level at a dose of 2.5 mg/kg

(25.34% decreases) as compared to diabetic control group table-3.

### Discussion

In the present study, Ethanolic extract of fruits of *Abelmoschus esculentus* at a dose of 500mg/kg b.w. could produce a significant fall in blood glucose levels by about 76% in diabetic rats, after 5 h of treatment. But none of these extracts could produce any hypoglycemic effect in normal rats. The aqueous extracts of fruits of *Abelmoschus esculentus* have not shown significant antihyperglycemic activity, hence the Ethanolic extracts may be considered to have good antihyperglycemic active principles without causing any hypoglycemic effect unlike insulin and other synthetic drugs. The phytochemical screening of fruits of *Abelmoschus esculentus* revealed the presence of alkaloids, carbohydrates, terpenoids, tannins, flavonoids and phenolic compounds and volatile oil. Flavonoids, alkaloids and phenolics are known to be bioactive antidiabetic principles. The antidiabetic effect of ethanolic extract of *Abelmoschus esculentus* fruits may be due to the presence of more than one antihyperglycemic principle and their synergistic properties. In this study, the antihyperglycemic activity caused by glibenclamide in alloxan-induced diabetic rats is an indication of the presence of some beta cells, as glibenclamide is known to stimulate insulin secretion from beta cells. The Ethanolic extract of *Abelmoschus esculentus* fruits may have stimulating effect on the remnant beta cells. However, further experiments are required to elucidate the exact mechanism of action. Further studies will be focused on determination of the mechanism(s) of action, as well as on the isolation of bioactive principles.

### Acknowledgements

The authors are grateful to Dr. Surendra Kumar Jain, director, Sagar Institute of Research and Technology,

Bhopal, India, for providing me the facility to carry out the pharmacological studies. We would like to thank Vtinerary College, Mahow, Indore; his help in carrying out the Histopathological parameters were studied.

### Reference

1. Valiathan MS. Healing plants. *Current Science*. 1998; 75 (11); 1122–26.
2. Larner J. The pharmacological basis of therapeutics. 7<sup>th</sup> ed. New York: Mac Millan; 1985.
3. The WHO Expert Committee on Diabetes Mellitus. Technical report series. Geneva: World Health Organization; 1980.
4. Manohar MS. Pod development and germination of bhindi (*Abelmoschus esculentus*). *Experimental Agriculture*. 1969; 5; 249-55.
5. Solanki SS, Singh RD, Yadav JP. Studies on the temperature and media relations and coefficient velocity of germination of vegetable seeds. II. Summer squash (*Cucurbita pepo* L.) and okra (*Abelmoschus esculentus* (L.) Moench.). *Progressive Horticulture*, 1980;12; 59-65.
6. Khomsug P, Thongjaroenbuangam W, Pakdeenarong N, Suttajit M, Chantiratikul P. Antioxidative activities and phenolic content of extracts from Okra (*Abelmoschus esculentus* L.) *Res J Biol Sci*. 2010;5:310–13
7. Bansal SP. Healing Power of Foods: Nature's Prescription of Common Diseases. Pustak Mahal Publishers, 2004; 32.
8. Sengkhamparn N, Verhoef R, Schols HA, Sajjaanantakul T, Voragen AG. Characterisation of cell wall polysaccharides from okra (*Abelmoschus esculentus* (L.) Moench) *Carbohydr Res*. 2009;344:1824–32.
9. Borssum Waalkes J V. *Malesian Malvaceae revised*. *Blumea*, 1966; 14; 100–101.

10. Tomoda M, Shimizu N, Gonda R, Kanari M, Yamada H, Hikino H. Anticomplementary and hypoglycemic activity of okra and hibiscus mucilage. *Carbohydr Res.* 1989;190:323–8.
11. Anonymous, Wealth of India-Raw materials H-K. Council of Scientific and Industrial Research, New Delhi, India. 1959; 84–8.
12. Siemonsma J S, Kasem Piluek, eds. Vegetables. Plant Resources of South-East Asia (PROSEA). (Pl Res SEAs) 1993; 8:57.
13. Feng Kuo-mei. Malvaceae. In: Feng Kuo-mei, ed., Fl. Reipubl. Popularis Sin. 1984; 49(2): 1-102.
14. Purseglove JW. Tropical Crops dicotyledons. Longman Group Limited. 1968; 368-70.
15. Siemonsma JS, Kouame C. *Abelmoschus Esculentus*. In plant resources of tropical Africa Vegetable. Editors Grubben G.J.H and O.A. Denton, Published by PROTA foundation Netherlands, 2004; 21-29.
16. Kochhar SL. Okra (Lady’s finger) In: Tropical crops, a textbook of economic Botany. Editor S.L., Kochhar, 1986; 263-64.
17. I-Min L, Shorong-Shii L, Ting-Wei L, Feng-Lin H, Juei-Tang C, Myricetin as the active principle of *Abelmoschus moschatus* to lower plasma glucose in streptozotocin-induced diabetic rats. *Planta Med.* 2005; 71 (7); 617-621.
18. Sathyanarayana U. In *Biochemistry*, 2nd edition, Upala Author – Publishers Interlinks, 2002; 587 – 592.
19. Kannur DM, Hukkeri VI. Antidiabetic activity of *Caesalpinia bonduella* seed extract in rats. *Fitoterapia.* 2006;10(1016), 546-549.
20. Mujumdar AM, Naik DG, Dandge CN, Puntambekar HM. Antiinflammatory activity of *Abelmoschus esculentus* Roxb. in albino rats. *Indian Journal of Pharmacology.* 2000; 32: 375-377.
21. Ghosh, SB, Gupta S, Chandra AK. Antifungal activity in fruits of *Abelmoschus esculentus* Roxb. *Indian J Exp Biol.* 1980 ;18(2):174-6.
22. Kameswara RB, Kesavulu MM, Giri R, Appa Rao, Ch., Antidiabetic and hypolipidemic effects of *Momordica cymbalaria*, Hookfruit powder in Alloxan diabetic rats. *Journal of Ethnopharmacology.* 1999; 67; 103–109.

**Table 1 Blood glucose level of aqueous and Ethanolic fruit extract of *Abelmoschus esculentus* during oral glucose tolerance test**

Groups (n=6)	Treatment and dose	Fasting Blood Glucose (mg/dl) (mean ± SEM)			
		0 min	30 min	60 min	120 min
Group 1	Normal control 10 ml/kg	89.33 ± 0.83	133.66 ± 0.56	117.33 ± 0.26	112.66 ± 1.38
Group 2	Standard 2.5 <sup>a</sup>	91.33 ± 0.80**	109.6 ± 0.50**	84.03 ± 0.16**	78.0 ± 0.97**
Group 3	Ethanolic extract 250 <sup>a</sup>	104.6 ± 0.46**	133.00 ± 0.63	116.66 ± 0.56	105.33 ± 0.35**
Group 4	Aqueous extract 250 <sup>a</sup>	99.33 ± 0.79**	127.0 ± 0.36**	114 ± 0.35**	101.66 ± 0.55**
Group 5	Ethanolic extract 500 <sup>a</sup>	101.6 ± 0.76**	129.0 ± 0.63**	116.33 ± 0.92	96.33 ± 0.91**
Group 6	Aqueous extract 500 <sup>a</sup>	104.0 ± 0.86**	125.6 ± 0.56**	115.0 ± 0.34*	102.66 ± 0.56**

a = mg/kg b.w.

Values ± SEM, n = no. of animal / group, [One way ANOVA followed by Dunnett test]\*\* P < 0.01, \* P < 0.05 v/s Normal control

**Table 2 Fasting blood glucose level of alloxan - induced diabetic rats in acute study**

Group (n = 6)	Treatment and dose	Fasting Blood Glucose (mg/dl)				
		0 hr	2 hr	4 hr	6 hr	24 hr
Group 1	Normal control 10 ml/kg	100.09 ± 0.43**	99.15 ± 1.06**	99.77 ± 1.04**	99.32 ± 1.07**	100.09 ± 1.12**
Group 2	Diabetic control 10 ml/kg	132.67 ± 1.27	135.33 ± 0.95	137.67 ± 2.5	140.0 ± 3.03	142.5 ± 3.33
Group 3	Standard 2.5 <sup>a</sup>	131.00 ± 1.61	114.17 ± 1.7**	106.33 ± 0.92**	100.83 ± 0.91**	97.0 ± 0.37**
Group 4	Ethanollic extract 250 <sup>a</sup>	131.05 ± 0.54	117.9 ± 0.24**	115.23 ± 0.71**	109.42 ± 0.91**	105.73 ± 0.26**
Group 5	Aqueous extract 250 <sup>a</sup>	129.45 ± 1.91	117.5 ± 1.56**	116.1 ± 0.11**	113.15 ± 0.23**	108.12 ± .81**
Group 6	Ethanollic extract 500 <sup>a</sup>	130.14 ± 1.4	117.0 ± 0.2**	113.23 ± 0.03**	113.13 ± 1.10**	100.33 ± 1.08**
Group 7	Aqueous extract 500 <sup>a</sup>	131.42 ± 1.81	121.0 ± 1.02**	115.7 ± 0.23**	112.12 ± 0.41**	104.12 ± 0.18**

a = mg/kg b.w.

Values ± SEM, n = no. of animal / group, [One way ANOVA followed by Dunnett test] \*\* P < 0.01 v/s Diabetic control

**Table 3 Fasting blood glucose level of alloxan - induced diabetic rats after 14 days subacute treatment**

Group (n = 6)	Treatment and dose	Fasting Blood Glucose (mg/dl)				
		1 <sup>st</sup> day	3 <sup>rd</sup> day	5 <sup>th</sup> day	10 <sup>th</sup> day	14 <sup>th</sup> day
Group 1	Normal control 10 ml/kg	101.0 ± 0.9**	102.6 ± 0.92**	101.6 ± 0.76**	102.3 ± 0.56**	100.7 ± 0.76**
Group 2	Diabetic control 10 ml/kg	133.17 ± 0.79	136.0 ± 1.34	139.67 ± 1.56	141.67 ± 1.41	145.33 ± 0.56
Group 3	Standard 2.5 <sup>a</sup>	130.83 ± 1.40	121.3 ± 1.65**	117.0 ± 1.93**	100.8 ± 1.40**	97.67 ± 0.56**
Group 4	Ethanollic extract 250 <sup>a</sup>	131.00 ± 1.34	128.0 ± 1.37**	122.7 ± 0.84**	121.3 ± 1.12**	114.6 ± 0.61**
Group 5	Aqueous extract 250 <sup>a</sup>	129.67 ± 1.38	127.3 ± 0.84**	125.3 ± 0.56**	122.0 ± 0.73**	115.7 ± 0.56**
Group 6	Ethanollic extract 500 <sup>a</sup>	132.33 ± 2.14	126.0 ± 0.73**	122.0 ± 0.73**	116.0 ± 1.77**	107.3 ± 3.16**
Group 7	Aqueous extract 500 <sup>a</sup>	130.67 ± 1.84	126.5 ± 1.12**	120 ± 0.73**	118.7 ± 0.56**	110.5 ± 1.23**

a = mg/kg b.w.

Values ± SEM, n = no. of animal / group, [One way ANOVA followed by Dunnett test]\*\* P < 0.01, \* P < 0.05 v/s Diabetic control